

FIRST AID POLICY

Last Reviewed/Updated: 01.05.2017

Next Review/Update: 01.05.2018

1. Statement of Commitment

Regent Independent College, hereafter referred to as 'Regent' is committed to caring for, and protecting, the health, safety and welfare of its students, staff and visitors. We confirm our adherence to the following standards at all times:

- To make practical arrangements for the provision of First Aid on our premises, during off-site sport and on school visits.
- To ensure that trained First Aid staff renew, update or extend their approved qualifications at least every three years.
- To have a minimum of 2 trained First Aiders on site at any one time, including a person with a paediatric first aid qualification. Such people will be able to responsibly deliver or organise emergency treatment.
- On any external trips, the accompanying staff member/s shall call for an ambulance for any kind of accident unless it is sufficiently minor that it can be dealt with using the first aid kit.
- To record accidents and illnesses appropriately, reporting to parents and the Health & Safety Executive under the Reporting of Injuries, Diseases and Dangerous Occurrences Regulations (1995).
- To provide accessible first aid kits at various locations on site, along with a portable kit for trips, excursions and sport.
- To record and make arrangements for students and staff with specific medical conditions.
- To deal with the disposal of bodily fluids and other medical waste accordingly, providing facilities for the hygienic and safe practice of first aid.
- To contact the medical emergency services if they are needed, informing next of kin immediately in such a situation.
- To communicate clearly to students and staff where they can find medical assistance if a person is ill or an accident has occurred.
- To communicate clearly in writing to parents or guardians if a child has sustained a bump to the head at school, however minor.

2.

Details of First Aid Practitioners at Regent

Appointed Person:	Qualification	Training Renewal Date
Vinita Gaur	Paediatric First Aider	8/12/19
Deboarah Yala	Paediatric First Aider	8/12/19
Lynne Bonfim	Paediatric First Aider	2/7/18
Caroline Shepheard	Paediatric First Aider	2/7/18

Practical Arrangements at Regent

Location of First Aid Facilities

- The sick room is located off the corridor opposite Reception for first aid treatment and for student or staff to rest/recover if feeling unwell. This includes; a bed, first aid supplies, a water supply and sink, an adjacent bathroom and hygiene supplies such as gloves and paper towels.
- A portable first aid kit must be obtained from the office for school visits.
- Other first aid kits are held at the following locations: in the first floor kitchen, one at reception and in the science lab

Responsibilities of the Trained First Aiders

- Provide appropriate care for students or staff who are ill or sustain an injury
- Record all accidents in the accident book (to be found in the Reception). They are then passed to the receptionist who will scan a copy onto the student's SIMS profile. In addition, the hard copy will be kept in a folder in the Exams Cupboard and will be held for three years before being disposed of.
- In the event of any injury to the head, however minor, ensure that notification is made to parents/guardians.
- Make arrangements with parents/guardians to collect students in the post 16 GCSE, Lower and Upper Sixth years and take them home if they are deemed too unwell to continue the school day, where possible, and in other cases, to ensure that a parent/guardian has been made aware that the student is unwell and that they are making their way home independently. For students in the senior school, that is, for pre16 GCSE, students may not leave the site before the school days ends until permission by a parent/guardian is given. However, it is deemed preferable that a parent/guardian collects the student from the site.

- The Co-Principal is to be informed in all cases where first aid has been given to a student.

The Co-Principal is to ensure that the following duties are delegated and carried out:

- Ensure that all staff and students are familiar with the school's first aid and medical procedures.
- Ensure that all staff are familiar with measures to provide appropriate care for students with particular medical needs (eg. Diabetic needs, Epi-pens, inhalers).
- Monitor and re-stock supplies and ensure that first aid kits are replenished.
- Ensure that the school has an adequate number of appropriately trained First Aiders.
- Co-ordinate First Aiders and arrange for training to be renewed as necessary.
- Maintain adequate facilities.
- Ensure that correct provision is made for students with special medical requirements both in school and on off-site visits.
- The first aid provision is monitored on a weekly basis.
- Fulfil the school's commitment to report to Reporting of Injuries, Diseases and Dangerous Occurrences Regulations (2013) - RIDDOR, as described below
- Liaise with managers of external facilities, such as the local sports facilities, to ensure appropriate first aid provision.
- Contact emergency medical services as required.
- Maintain an up-to-date knowledge and understanding of guidance and advice from appropriate agencies

What to do in the case of an accident, injury or illness

A member of staff or student witnessing an accident, injury or illness should immediately contact a named trained first aider (see above). The school office should be contacted if the location of a trained first aider is uncertain. Any student or member of staff sustaining an injury whilst at school should be seen by a first aider who will provide immediate first aid and summon additional help as needed. The student or member of staff should not be left unattended. The first aider will organise an injured student's transfer to the sick room if possible and appropriate and to hospital in the case of an emergency. Parents should be informed as necessary by telephone by the first aider or receptionist. A written record of all accidents and injuries is maintained in the accident book.

Contacting parents

Parents should be informed by telephone as soon as possible after an emergency or following a **serious/significant** injury including:

- Head injury
- Suspected sprain or fracture

- Following a fall from height
- Dental injury
- Anaphylaxis & following the administration of an Epi-pen
- Epileptic seizure
- Severe hypoglycaemia for students, staff or visitors with diabetes
- Severe asthma attack
- Difficulty breathing
- Bleeding injury
- Loss of consciousness
- If the student is generally unwell

If non-emergency transportation is required, an authorised taxi service will be used if parents are delayed. A member of staff will accompany the student until a parent arrives.

Contacting the Emergency Services

An ambulance should be called for any condition listed above or for any injury that requires emergency treatment. Any student taken to hospital by ambulance must be accompanied by a member of staff until a parent arrives. All cases of a student becoming unconsciousness (not including a faint) or following the administration of an Epi-pen, must be taken to hospital.

Accident reporting

The accident book must be completed for any accident or injury occurring onsite, at the local sports facilities, or on a school trip. This includes any accident involving staff or visitors. The accident book will be monitored by the Head Teacher as certain injuries require reporting (RIDDOR requirements).

Students who are unwell in school

Any student who is unwell cannot be left to rest unsupervised in the sick room. If a student becomes unwell, a parent should be contacted as soon as possible by a First Aider, a member of the school admin team or the Principal. Anyone not well enough to be in school should be collected as soon as possible by a parent. Staff should ensure when a student goes home it is noted as such by the person keeping the school registers.

First Aid equipment and materials

The Principal will delegate responsibility for stocking and checking the first aid kits to a senior academic administrator. The first aid boxes contain:

- A first aid guidance card
- At least 20 adhesive hypo allergenic plasters (including blue plasters for home economics)
- 4 triangular bandages (slings)

- Safety pins
- Cleaning wipes
- Adhesive tape
- 2 sterile eye pads
- 6 medium sized unmedicated dressings
- 2 large sized unmedicated dressings
- Disposable gloves
- 1 resuscitator
- Yellow clinical waste bag

First aid for school trips

The trip organiser must undertake a risk assessment to ensure an appropriate level of first aid cover, with reference to the Educational Visits Policy, which includes further guidance. A First Aid kit for school trips must be collected from the main office. This must be returned to the main office for replenishing on return. Any accidents/injuries must be reported to the appointed person and to parents and documented in the accident book in accordance with this policy. RIDDOR guidelines for reporting accidents must be adhered to. For any major accident or injury the appropriate health & safety procedure must be followed.

Students using crutches or having limited mobility

Parents must inform the school of the nature of injury and the anticipated duration of immobility. Information about the condition will be disseminated to teachers so they are fully aware of the student's needs. Arrangements will be made for the student to arrive/leave lessons early to allow for a safe transfer around school. Parents must inform the school of any particular difficulties.

Students with medical conditions

A list of students with pre-existing medical conditions will be kept by the Co-Principal. This information is useful for lesson planning and for risk assessments prior to a school trip.

Dealing with body fluids

In order to maintain protection from disease, all body fluids should be considered infected. To prevent contact with body fluids. Bodily fluids include Blood, faeces, nasal and eye discharges, saliva and vomit. When dealing with bodily fluids, the following guidelines should be followed.

- When dealing with any body fluids wear disposable gloves.
- Wash hands thoroughly with soap and warm water after the incident.
- Keep any abrasions covered with a plaster.
- Spills of the following body fluids must be cleaned up immediately.

Disposable towels should be used to soak up the excess, and then the area should be treated with a disinfectant solution. Never use a mop for cleaning up blood and body fluid spillages. All contaminated material should be disposed of in a yellow clinical waste bag (available in all 1st aid boxes) then placed in the waste bin in the sick room. Avoid getting any body fluids in your eyes, nose, and mouth or on any open sores. If a splash occurs, wash the area well with soap and water or irrigate with copious amounts of saline.

Infectious diseases

If a student is suspected of having an infectious disease advice should be sought from the Co-Principal who will follow the Health Protection Agency guidelines below to reduce the transmission of infectious diseases to other students and staff.

ILLNESS	PERIOD OF EXCLUSION	COMMENTS
Chickenpox	5 days from onset of rash	Pregnant women up to 20 weeks and those in last 3 weeks of pregnancy should inform their midwife that they have been in contact with chickenpox. Any children being treated for cancer or on high doses of steroids should also seek medical advice.
German Measles	For 5 days from onset of rash	Pregnant women should inform their midwife about contact
Impetigo	Until lesions are crusted or healed	Antibiotic treatment by mouth may speed healing
Measles	5 days from onset of rash	Any children being treated for cancer or on high doses of steroids must seek medical advice
Scabies	Until treatment has been commenced	Two treatments one week apart for cases. Treatment should include all household members and any other very close contacts
Scarlet Fever	5 days after commencing antibiotics	Antibiotic treatment recommended
Slapped Cheek Syndrome	None	Pregnant women up to 20 weeks must inform their midwife about contact
Diarrhoea and vomiting	48 hours from last episode of diarrhoea or vomiting	Exclusion from swimming may be needed
Hepatitis A	Exclusion may be necessary	Consult the Health Protection Agency
Meningococcal meningitis	Until recovered	Communicable disease control will give advice on any treatment needed and identify contact requiring treatment. No need to exclude siblings or other close contacts.
Viral Meningitis	Until fully recovered	Milder illness
Threadworms	None	Treatment is recommended for the student and family members
Mumps	5 days from onset of swollen glands	
Head Lice	None once treated	Treatment is recommended for the student and close contacts if live lice are found
Conjunctivitis	None	Children do not usually need to stay off school with conjunctivitis if they are feeling well. If, however, they are feeling unwell with conjunctivitis they should stay off school until they feel better
Influenza	Until fully recovered	
Cold sores	None	Avoid contact with the sores
Warts, verrucae	None	Verrucae should be covered in swimming pools, gymnasiums and changing rooms
Glandular fever	None	
Tonsillitis	None	

Medication in School

The school aims to support as far as possible, and maintain the safety of, students who require medication during the school day.

However, it should be noted that:

- No child should be given any medication without their parent's written consent.
- No Aspirin products are to be given to any student at school.
- Parents must be given written confirmation of any medication administered at school, a copy of which will be filed on SIMS.

Students who need to take medication during the school day e.g. antibiotics. However, wherever possible the timing and dosage should be arranged so that the medication can be administered at home.

(i) Non-Prescription Medication

These are only to be administered by the appointed person or a designated person if they have agreed to this extension of their role and have been appropriately trained.

A teacher may administer non-prescription medication on a residential school trip provided that written consent* has been obtained in advance. This may include travel sickness pills or pain relief.

All medication administered must be documented, signed for and parents informed in writing.

* Parents are asked to complete a consent form at the start of the academic year to cover the administration of non-prescription medicines when deemed necessary by a school first aider. Under changes to Early Years Foundation Stage (EYFS) guidance (January 2012), this is now also acceptable in the case for EYFS students, provided that parents are contacted immediately before the administration of the medication. In all cases which rely on such on-going consent, parents must, nevertheless, be informed in writing that the administration of medication has taken place.

(ii) Prescription-Only Medication

Prescribed medicines may be given to a student by the appointed person or a designated person if they have agreed to this extension of their role and have been appropriately trained.

Written consent must be obtained from the parent or guardian, clearly stating the name of the medication, dose, frequency and length of course.

The school will accept medication from parents only if it is in its original container

A form for the administration of medicines in school is available from the appointed person.

(iii) Administration of Medication

- The medication must be checked before administration by the member of staff confirming the medication name, student name, dose, time to be administered and the expiry date.
- Wash hands.
- Confirm that the student's name matches the name on the medication

- Explain to the student that his or her parents have requested the administration of the medication.
- Document, date and sign for what has been administered.
- Complete the form which goes back to parents
- Ensure that the medication is correctly stored in a locked drawer or cupboard, out of the reach of students.
- Antibiotics and any other medication which requires refrigeration should be stored in the fridge in the staff kitchen on the Ground Floor. All medication should be clearly labelled with the student's name and dosage.
- Parents should be asked to dispose of any out of date medication.
- Used needles and syringes must be disposed of in the sharps box kept in the sick room.

(iv) Emergency Medication

It is the parents' responsibility to inform the school of any long-term medical condition that may require regular or emergency medication to be given. In these circumstances a health care plan may be required and this will be completed and agreed with parents.

Guidelines for reporting: RIDDOR (Reporting of Injuries, Diseases and Dangerous Occurrences Regulations 10/13)

By law any of the following accidents or injuries to students, staff, visitors, members of the public or other people not at work requires notification to be sent to the Health and Safety executive by phone, fax, email or letter.

Major injuries from schedule 1 of the regulations:

1. Any fracture, other than to the fingers, thumbs or toes.
2. Any amputation.
3. Dislocation of the shoulder, hip, knee or spine.
4. Loss of sight (whether temporary or permanent)
5. A chemical or hot metal burn to the eye or any penetrating injury to the eye.
6. Any injury resulting from an electric shock or electrical burn (including any electrical burn caused by arcing or arcing products, leading to unconsciousness or requiring resuscitation or admittance to hospital for more than 24 hours.
7. Any other injury leading to hypothermia, heat induced illness or to unconsciousness requiring resuscitation or admittance to hospital for more than 24 hours
8. Any other injury lasting over 3 days
9. Loss of consciousness caused by asphyxia or by exposure to a harmful substance or biological agent.
10. Either of the following conditions which result from the absorption of any substance by inhalation, ingestion or through the skin:
 - Acute illness requiring medical treatment; or

- Loss of consciousness
11. Acute illness which requires medical treatment where there is reason to believe that this resulted from exposure to a biological agent or its toxins or infected material.
 12. Death
 13. A specified dangerous occurrence, where something happened which did not result in an injury, but could have done.

A copy of this policy is available on the College website and also in administration R-drive under school policies.

APPENDIX: Guidance to staff on particular medical conditions

(i) Allergic reactions

Symptoms and treatment of a mild allergic reaction:

- Rash
- Flushing of the skin
- Itching or irritation

If the student has a care plan, follow the guidance provided and agreed by parents. Administer the prescribed dose of antihistamine to a child who displays these mild symptoms only. Make a note of the type of medication, dose given, date, and time the medication was administered. Complete and sign the appropriate medication forms, as detailed in the policy. Observe the child closely for 30 minutes to ensure symptoms subside.

(ii) Anaphylaxis

Symptoms and treatment of Anaphylaxis:

- Swollen lips, tongue, throat or face
- Nettle type rash
- Difficulty swallowing and/or a feeling of a lump in the throat
- Abdominal cramps, nausea and vomiting
- Generalised flushing of the skin
- Difficulty in breathing
- Difficulty speaking
- Sudden feeling of weakness caused by a fall in blood pressure
- Collapse and unconsciousness

When someone develops an anaphylactic reaction the onset is usually sudden, with the following signs and symptoms of the reaction progressing rapidly, usually within a few minutes.

Action to be taken

1. Send someone to call for a paramedic ambulance and inform parents. Arrange to meet parents at the hospital.
2. Send for the named emergency box.
3. Reassure the student help is on the way.
4. Remove the Epi-pen from the carton and pull off the grey safety cap.
5. Place the black tip on the student's thigh at right angles to the leg (there is no need to remove clothing).
6. Press hard into the thigh until the auto injector mechanism functions and hold in place for 10 seconds.
7. Remove the Epi-pen from the thigh and note the time.

8. Massage the injection area for several seconds.
9. If the student has collapsed lay him/her on the side in the recovery position.
10. Ensure the paramedic ambulance has been called.
11. Stay with the student.
12. Steps 4-8 maybe repeated if no improvement in 5 minutes with a second Epi-pen if you have been instructed to do so by a doctor.

REMEMBER Epi-pens are not a substitute for medical attention, if an anaphylactic reaction occurs and you administer the Epi-pen the student must be taken to hospital for further checks.

Epi-pen treatment must only be undertaken by staff who have received specific training.

(iii) Asthma management

The school recognises that asthma is a serious but controllable condition and the school welcomes any student with asthma. The school ensures that all students with asthma can and do fully participate in all aspects of school life, including any out of school activities. Taking part in PE is an important part of school life for all students and students with asthma are encouraged to participate fully in all PE lessons. Teaching staff will be aware of any child with asthma from a list of students with medical conditions kept in the staff room. The school has a smoke free policy.

Trigger factors

- Change in weather conditions
- Animal fur
- Having a cold or chest infection
- Exercise
- Pollen
- Chemicals
- Air pollutants
- Emotional situations
- Excitement

General considerations

Students with asthma need immediate access to their reliever inhaler. Younger students will require assistance to administer their inhaler. It is the parents' responsibility to ensure that the school is provided with a named, in-date reliever inhaler, which is kept in the classroom, not locked away and always accessible to the student. Teaching staff should be aware of a child's trigger factors and try to avoid any situation that may cause a student to have an asthma attack. It is the parents' responsibility to provide a new inhaler when out of date. Students must be made aware of where their inhaler is kept and this medication must be taken on any out of school activities.

As appropriate for their age and maturity, students are encouraged to be responsible for their reliever inhaler, which is to be brought to school and kept in a school bag to be used as required.

A spare named inhaler should be brought to school and given to the class teacher for use if the student's inhaler is lost or forgotten.

Recognising an asthma attack

- Student unable to continue an activity
- Difficulty in breathing
- Chest may feel tight
- Possible wheeze
- Difficulty speaking
- Increased anxiety
- Coughing, sometimes persistently

Action to be taken

1. Ensure that prescribed reliever medication (usually blue) is taken promptly.
2. Reassure the student.
3. Encourage the student to adopt a position which is best for them-usually sitting upright.
4. Wait five minutes. If symptoms disappear the student can resume normal activities.
5. If symptoms have improved but not completely disappeared, inform parents and give another dose of their inhaler and call the appointed person or a first aider if she not available.
6. Loosen any tight clothing.
7. If there is no improvement in 5-10 minutes continue to make sure the student takes one puff of their reliever inhaler every minute for five minutes or until symptoms improve.
8. Call an ambulance.
9. Accompany student to hospital and await the arrival of a parent.

(iv) Diabetes management

Students with diabetes can attend school and carry out the same activities as their peers but some forward planning may be necessary. Staff must be made aware of any student with diabetes attending school.

Signs and symptoms of low blood sugar (hypoglycaemic attack)

This happens very quickly and may be caused by: a late meal, missing snacks, insufficient carbohydrate, more exercise, warm weather, too much insulin and stress. The student should test his or her blood glucose levels if blood testing equipment is available.

- Pale
- Glazed eyes
- Blurred vision
- Confusion/incoherent
- Shaking
- Headache
- Change in normal behaviour-weepy/aggressive/quiet

- Agitated/drowsy/anxious
- Tingling lips
- Sweating
- Hunger
- Dizzy

Action to be taken

1. Follow the guidance provided in the care plan agreed by parents.
2. Give fast acting glucose-either 50ml glass of Lucozade or 3 glucose tablets. (Students should always have their glucose supplies with them. Extra supplies will be kept in emergency boxes. This will raise the blood sugar level quickly.
3. This must be followed after 5-10 minutes by 2 biscuits, a sandwich or a glass of milk.
4. Do not send the child out of your care for treatment alone.
5. Allow the student to have access to regular snacks.
6. Inform parents.

Action to take if the student becomes unconscious:

1. Place student in the recovery position and seek the help of the appointed person or a first aider.
2. Do not attempt to give glucose via mouth as student may choke.
3. Telephone 999.
4. Inform parents.
5. Accompany student to hospital and await the arrival of a parent.

Signs and symptoms of high blood sugar (hyperglycaemic attack)

Hyperglycaemia – develops much more slowly than hypoglycaemia but can be more serious if left untreated. It can be caused by too little insulin, eating more carbohydrate, infection, stress and less exercise than normal.

- Feeling tired and weak
- Thirst
- Passing urine more often
- Nausea and vomiting
- Drowsy
- Breath smelling of acetone
- Blurred vision
- Unconsciousness

Action to be taken

1. Inform the appointed person or a first aider
2. Inform parents
3. Student to test blood or urine

4. Call 999

(v) Epilepsy management

How to recognise a seizure

There are several types of epilepsy but seizures are usually recognisable by the following symptoms:

- Student may appear confused and fall to the ground.
- Slow noisy breathing.
- Possible blue colouring around the mouth returning to normal as breathing returns to normal.
- Rigid muscle spasms.
- Twitching of one or more limbs or face
- Possible incontinence.

A student diagnosed with epilepsy will have an emergency care plan.

Action to be taken

1. Send for an ambulance;
 - a. if this is a student's first seizure,
 - b. if a student known to have epilepsy has a seizure lasting for more than five minutes or
 - c. if an injury occurs.
2. Seek the help of the appointed person or a first aider.
3. Help the student to the floor.
4. Do not try to stop seizure.
5. Do not put anything into the mouth of the student.
6. Move any other students away and maintain student's dignity.
7. Protect the student from any danger.
8. As the seizure subsides, gently place them in the recovery position to maintain the airway.
9. Allow patient to rest as necessary.
10. Inform parents.
11. Call 999 if you are concerned.
12. Describe the event and its duration to the paramedic team on arrival.
13. Reassure other students and staff.
14. Accompany student to hospital and await the arrival of a parent.

